

病的知覚体験と他者の表象

Pathological perceptual experiences and representations of others

西尾 慶之
Yoshiyuki Nishio

東京都立松沢病院
Tokyo Metropolitan Matsuzawa Hospital
nishiokov@gmail.com

Abstract

A variety of pathological perceptual symptoms, including visual hallucinations of persons, a sense of presence and kinetopsia, are observed in patients with dementia with Lewy bodies and Parkinson's disease. I propose that these symptoms may develop with a series of psychological steps, in which aberrant spatial processes leads to presence feelings of illusory/hallucinatory objects, and then activate representations of others. I will present phenomenological and neurological evidence that supports this idea.

Keywords — **noisy perception, abnormal cortical state, delusional acceptance**

レビー小体型認知症やパーキンソン病では、人の幻視、実体意識性（「近くに誰かがいる」という幻覚）、キネトプシア（動きの錯視）などの病的知覚体験が観察される。演者は「逸脱した空間知覚→幻の対象の存在感→他者にまつわる意味表象の賦活」という心理の流れに基づいてこれらの病的知覚体験を理解できるのではないかと考えている。本発表では、この仮説の根拠となる現象的、神経学的知見について紹介する。

参考文献

- [1] Nishio Y, Yokoi K, Hirayama K, Ishioka T, Hosokai Y, Gang M, Uchiyama M, Baba T, Suzuki K, Takeda A, Mori E. Defining visual illusions in Parkinson's disease: Kinetopsia and object misidentification illusions. *Parkinsonism Relat Disord.*2018. doi: 10.1016/j.parkreldis.2018.05.023. [Epub ahead of print]
- [2] Watanabe H, Nishio Y, Mamiya Y, Narita W, Iizuka O, Baba T, Takeda A, Shimomura T, Mori E. Negative mood invites psychotic false perception in dementia. *PLoS One.* 2018;13(6):e0197968. doi: 10.1371/journal.pone.0197968.
- [3] Nishio Y, Yokoi K, Uchiyama M, Mamiya Y, Watanabe H, Gang M, Baba T, Takeda A, Hirayama K, Mori E. Deconstructing psychosis and misperception symptoms in Parkinson's disease. *J Neurol Neurosurg Psychiatry.* 2017;88(9):722-729. doi: 10.1136/jnnp-2017-315741.